

BILINEAR SPACE-TIME ESTIMATES FOR LINEARISED KP-TYPE EQUATIONS ON THE THREE-DIMENSIONAL TORUS WITH APPLICATIONS

AXEL GRÜNROCK

ABSTRACT. A bilinear estimate in terms of Bourgain spaces associated with a linearised Kadomtsev-Petviashvili-type equation on the three-dimensional torus is shown. As a consequence, time localized linear and bilinear space time estimates for this equation are obtained. Applications to the local and global well-posedness of dispersion generalised KP-II equations are discussed. Especially it is proved that the periodic boundary value problem for the original KP-II equation is locally well-posed for data in the anisotropic Sobolev spaces $H_x^s H_y^\varepsilon(\mathbb{T}^3)$, if $s \geq \frac{1}{2}$ and $\varepsilon > 0$.

1. INTRODUCTION AND MAIN RESULTS

In a recent paper [7] joint with M. Panthee and J. Silva we investigated local and global well-posedness issues of the Cauchy problem for the dispersion generalised Kadomtsev-Petviashvili-II (KP-II) equation

$$(1) \quad \begin{cases} \partial_t u - |D_x|^\alpha \partial_x u + \partial_x^{-1} \Delta_y u + u \partial_x u = 0 \\ u(0, x, y) = u_0(x, y) \end{cases}$$

on the cylinders $\mathbb{T} \times \mathbb{R}$ and $\mathbb{T} \times \mathbb{R}^2$, respectively. We considered data u_0 satisfying the mean zero condition

$$(2) \quad \int_0^{2\pi} u_0(x, y) dx = 0$$

and belonging to the anisotropic Sobolev spaces $H_x^s(\mathbb{T})H_y^\varepsilon(\mathbb{R}^{n-1})$, $n \in \{2, 3\}$. We could prove quite general (with respect to the dispersion parameter α) local well-posedness results, to a large extent optimal - up to the endpoint - (with respect to the Sobolev regularity). In two dimensions and for higher dispersion ($\alpha > 3$) in three dimensions, these local results could be combined with the conservation of the L^2 -norm to obtain global well-posedness.

A key tool to obtain these results were certain bilinear space time estimates for free solutions, similar to Strichartz estimates. A central argument to obtain the space time estimates was the following simple observation. Consider a linearised version of (1) with a more general phase function

$$(3) \quad \begin{cases} \partial_t u - i\phi(D_x, D_y)u := \partial_t u - i\phi_0(D_x)u + \partial_x^{-1} \Delta_y u = 0 \\ u(0, x, y) = u_0(x, y) \end{cases}$$

1991 *Mathematics Subject Classification.* 35Q53.

The author was partially supported by the Deutsche Forschungsgemeinschaft, Sonderforschungsbereich 611.

where ϕ_0 is arbitrary at the moment, with solution $u(x, y, t) = e^{it\phi(D_x, D_y)}u_0(x, y)$. Then we can take the partial Fourier transform $\mathcal{F}_x$ with respect to the first spatial variable x only to obtain

$$\mathcal{F}_x e^{it\phi(D_x, D_y)}u_0(k, y) = e^{it\phi_0(k)}e^{i\frac{t}{k}\Delta_y}\mathcal{F}_x u_0(k, y).$$

Fixing k we have a solution of the free Schrödinger equation - with rescaled time variable $s := \frac{t}{k}$, and multiplied by a phase factor of size one. Now the whole Schrödinger theory - Strichartz estimates, bilinear refinements thereof, local smoothing and maximal function estimates - is applicable to obtain space time estimates for the linearised KP-type equation (3).

While in two space dimensions this simple argument has to be supplemented by further estimates depending on ϕ_0 , we could obtain (almost) sharp estimates in the three-dimensional $\mathbb{T} \times \mathbb{R}^2$ -case *only* by using the "Schrödinger trick" described above. In view of Bourgain's L^4_{xt} -estimate for free solutions of the Schrödinger equation with data defined on the two-dimensional torus [3, first part of Prop. 3.6], the question comes up naturally, if our analysis in [7] concerning $\mathbb{T} \times \mathbb{R}^2$ can be extended to KP-type equations on $\mathbb{T}^3$, and that's precisely the aim of the present paper.

To state our main results we have to introduce some more notation: We will consider functions $u, v, \dots$ of $(x, y, t) \in \mathbb{T} \times \mathbb{T}^2 \times \mathbb{R}$ with Fourier transform $\widehat{u}, \widehat{v}, \dots$, sometimes written as $\mathcal{F}u, \mathcal{F}v, \dots$, depending on the dual variables $(\xi, \tau) := (k, \eta, \tau) \in \mathbb{Z} \times \mathbb{Z}^2 \times \mathbb{R}$. Throughout the paper we assume $u, v, \dots$ to fulfill the mean zero condition $\widehat{u}(0, \eta, \tau) = 0$. For these functions we define the norms

$$\|u\|_{X_{s, \varepsilon, b}} := \| |k|^s \langle \eta \rangle^\varepsilon \langle \sigma \rangle^b \widehat{u} \|_{L^2_{\xi, \tau}},$$

where $\langle x \rangle^2 = 1 + |x|^2$ and $\sigma = \tau - \phi(\xi) = \tau - \phi_0(k) + \frac{|\eta|^2}{k}$. Although some of our arguments do not rely on that, we will always assume ϕ_0 to be odd, in order to have $\|u\|_{X_{s, \varepsilon, b}} = \|\overline{u}\|_{X_{s, \varepsilon, b}}$. For $\varepsilon = 0$ we abbreviate $\|u\|_{X_{s, \varepsilon, b}} = \|u\|_{X_{s, b}}$. In these terms our central bilinear space time estimate reads as follows.

Theorem 1. *Let $b > \frac{1}{2}$, $s_{1,2} \geq 0$ with $s_1 + s_2 > 1$ and $\varepsilon_{0,1,2} \geq 0$ with $\varepsilon_0 + \varepsilon_1 + \varepsilon_2 > 0$. Then the estimate*

$$(4) \quad \|D_y^{-\varepsilon_0}(uv)\|_{L^2_{xyt}} \lesssim \|u\|_{X_{s_1, \varepsilon_1, b}} \|v\|_{X_{s_2, \varepsilon_2, b}}$$

and its dualized version

$$(5) \quad \|uv\|_{X_{-s_1, -\varepsilon_1, -b}} \lesssim \|D_y^{\varepsilon_0}u\|_{L^2_{xyt}} \|v\|_{X_{s_2, \varepsilon_2, b}}$$

hold true.

Taking $\varepsilon_0 = 0$ and $u = v$ we obtain the linear estimate

$$(6) \quad \|u\|_{L^4_{xyt}} \lesssim \|u\|_{X_{s, \varepsilon, b}},$$

whenever $s, b > \frac{1}{2}$ and $\varepsilon > 0$. The estimate (4) can be applied to time localised solutions $e^{it\phi(D_x, D_y)}u_0$ and $e^{it\phi(D_x, D_y)}v_0$ of (3) to obtain

$$(7) \quad \|D_y^{-\varepsilon_0}(e^{it\phi(D_x, D_y)}u_0 e^{it\phi(D_x, D_y)}v_0)\|_{L^2_t([0,1], L^2_{xy})} \lesssim \|u_0\|_{H_x^{s_1} H_y^{\varepsilon_1}} \|v_0\|_{H_x^{s_2} H_y^{\varepsilon_2}},$$

provided $s_{1,2}$ and $\varepsilon_{0,1,2}$ fulfill the assumptions in Theorem 1. Especially for $s > \frac{1}{2}$ and $\varepsilon > 0$ we have the linear estimate

$$(8) \quad \|e^{it\phi(D_x, D_y)}u_0\|_{L^4_t([0,1], L^4_{xy})} \lesssim \|u_0\|_{H_x^s H_y^\varepsilon}.$$

The corresponding estimate for data u_0 defined on $\mathbb{R}^3$ holds global in time and has a $\|u_0\|_{H_x^{\frac{1}{2}} L_y^2}$ on the right hand side. It goes back to Ben Artzi and Saut [1].

Dimensional analysis shows that the Sobolev exponent $s = \frac{1}{2}$ is necessary. So we haven't lost more than an ε derivative in the x - as well as in the y -variable.

In order to prove Theorem 1, we will work in Fourier space, where the product uv is turned into the convolution

$$\widehat{u} * \widehat{v}(\xi, \tau) = \int d\tau_1 \sum_{\xi_1 \in \mathbb{Z}^3} \widehat{u}(\xi_1, \tau_1) \widehat{v}(\xi_2, \tau_2).$$

Here always $(\xi, \tau) = (k, \eta, \tau) = (k_1 + k_2, \eta_1 + \eta_2, \tau_1 + \tau_2) = (\xi_1 + \xi_2, \tau_1 + \tau_2)$. Observe that there is no contribution to the above sum, whenever $k_1 = 0$ or $k_2 = 0$. In the estimation of such convolutions the σ -weights in the $X_{s,\varepsilon,b}$ -norms become $\sigma_1 = \tau_1 - \phi(\xi_1)$ and $\sigma_2 = \tau_2 - \phi(\xi_2)$. With this notation we introduce the bilinear Fourier multiplier $M^{-\varepsilon}$, which we define by

$$\mathcal{F}M^{-\varepsilon}(u, v)(\xi, \tau) := \chi_{\{k \neq 0\}} \int d\tau_1 \sum_{\xi_1 \in \mathbb{Z}^3} \langle k_1 \eta - k \eta_1 \rangle^{-\varepsilon} \widehat{u}(\xi_1, \tau_1) \widehat{v}(\xi_2, \tau_2).$$

Observe that $|k_1 \eta - k \eta_1| = |k_1 \eta_2 - k_2 \eta_1|$, so that we have symmetry between u and v . The operator $M^{-\varepsilon}$ serves to compensate for the unavoidable loss of the D_y^ε in (4). A careful examination of the proof of Theorem 1 will give the following.

Theorem 2. *Let $s, b > \frac{1}{2}$ and $\varepsilon > 0$. Then*

$$(9) \quad \|M^{-\varepsilon}(u, v)\|_{L_{xyt}^2} \lesssim \|u\|_{X_{s,b}} \|v\|_{X_{s,b}}.$$

The proof of the above theorems will be done in section 2, while section 3 is devoted to the applications. Here we specialize to the dispersion generalised KP-II equation (1), that is to $\phi_0(k) = |k|^\alpha k$. For $\alpha = 2$, which is the original KP-II equation we will use Theorem 1 to show the following local result.

Theorem 3. *Let $s \geq \frac{1}{2}$ and $\varepsilon > 0$. Then, for $\alpha = 2$, the Cauchy problem (1) is locally well-posed for data $u_0 \in H_x^s H_y^\varepsilon(\mathbb{T}^3)$ satisfying the mean zero condition (2).*

For high dispersion, i. e. $\alpha > 3$, one can allow $s < 0$ and $\varepsilon = 0$. In fact, by the aid of Theorem 2 we can prove:

Theorem 4. *Let $3 < \alpha \leq 4$ and $s > \frac{3-\alpha}{2}$. Then the Cauchy problem (1) is locally well-posed for data $u_0 \in H_x^s L_y^2(\mathbb{T}^3)$ satisfying (2). If $s \geq 0$ the corresponding solutions extend globally in time by the conservation of the L_{xy}^2 -norm.*

More precise statements of the last two theorems will be given in section 3. We conclude this introduction with several remarks commenting on our well-posedness results and their context.

1. Concerning the Cauchy problem for the KP-II equation and its dispersion generalisations on $\mathbb{R}^2$ and $\mathbb{R}^3$ there is a rich literature, see e. g. [8], [9], [12], [13], [16], [17], [20], [22], this list is by no means exhaustive. For $\alpha = 2$ the theory has even been pushed to the critical space in a recent work of Hadac, Herr, and Koch [10]. On the other hand, for the periodic or semiperiodic problem the theory is much less developed. Besides Bourgain's seminal paper [2] our only references here are the papers [18], [19] of Saut and Tzvetkov and our own contribution [7] joint with M. Panthee and J. Silva.

2. The results obtained here for the fully periodic case are as good as those in [7] for the $\mathbb{T} \times \mathbb{R}^2$ case and even as those obtained by Hadac [8] for $\mathbb{R}^3$, which are optimal by scaling considerations. We believe this is remarkable since apart from

nonlinear wave and Klein-Gordon equations there are only very few examples in the literature, where the periodic problem is as well behaved as the corresponding continuous case. (One example is of course Bourgain's $L_x^2(\mathbb{T})$ result for the cubic Schrödinger equation [3], but this is half a derivative away from the scaling limit.) On the other hand there are many examples, such as KdV and mKdV, where at least the methods applied here lead to (by $\frac{1}{4}$ derivative) weaker results for the periodic problem. Another example is the KP-II equation itself in two space dimensions, where in [7] we lost $\frac{1}{4}$ derivative when stepping from $\mathbb{R}^2$ to $\mathbb{T} \times \mathbb{R}$. Another loss of $\frac{1}{4}$ derivative in the step from $\mathbb{T} \times \mathbb{R}$ to $\mathbb{T}^2$ is probable.

3. For the semilinear Schrödinger equation

$$iu_t + \Delta u = |u|^p u$$

on the torus, with $2 < p < 4$ in one, $1 < p < 2$ in two dimensions, one barely misses the conserved L_x^2 norm and thus cannot infer global well-posedness. The reason behind that is the loss of an ε derivative in the Strichartz type estimates in the periodic case. A corresponding derivative loss is apparent in Theorem 1 but the usually ignored mixed part of the rather comfortable resonance relation of the dispersion generalised KP-II equation allows (via $M^{-\varepsilon}$) to compensate for this loss, so that for high dispersion ($3 < \alpha \leq 4$) we can obtain something global. The author did not expect that, when starting this investigation.

4. We restrict ourselves to the most important (as we believe) values of α . Our arguments work as well for $\alpha \in (2, 3]$ with optimal lower bound for s but possibly with an ε loss in the y variable. For $\alpha > 4$ we probably lose optimality.

5. In [21] Takaoka and Tzvetkov proved a time localised $L^4 - L^2$ Strichartz type estimate *without* derivative loss for free solutions of the Schrödinger equation with data defined on $\mathbb{R} \times \mathbb{T}$. Inserting their arguments in our proof of Theorem 1 we can show a variant thereof with $\varepsilon_0 = \varepsilon_1 = \varepsilon_2 = 0$, if the data live on $\mathbb{T} \times \mathbb{R} \times \mathbb{T}$. Consequently our well-posedness results are valid in this case, too.

2. PROOF OF THEOREM 1

The main ingredient in the proof of Bourgain's Schrödinger estimate

$$\|e^{it\Delta} u_0\|_{L_{xt}^4(\mathbb{T}^3)} \lesssim \|u_0\|_{H_x^\varepsilon(\mathbb{T}^2)}, \quad (\varepsilon > 0)$$

is the well known estimate on the number of representations of an integer $r > 0$ as a sum of two squares: For any $\varepsilon > 0$ there exists c_ε such that

$$(10) \quad \#\{\eta \in \mathbb{Z}^2 : |\eta|^2 = r\} \leq c_\varepsilon r^\varepsilon.$$

For (10), see [11, Theorem 338]. Our proof of Theorem 1 relies on the following variant thereof.

Lemma 1. *Let $r \in \mathbb{N}$, $\delta \in \mathbb{R}^2$. Then for any $\varepsilon > 0$ there exists c_ε , independent of r and δ , such that*

$$(11) \quad \#\{\eta \in \mathbb{Z}^2 : r \leq |\eta - \delta|^2 < r + 1\} \leq c_\varepsilon r^\varepsilon.$$

Proof. In the case where $\delta \in \mathbb{Z}^2$, this follows by translation from (10). So we may assume $\delta \in [0, 1]^2$, and we start by considering the special case $\delta = (\frac{1}{2}, \frac{1}{2})$. Here

$$\begin{aligned} & \#\{\eta \in \mathbb{Z}^2 : r \leq |\eta - \delta|^2 < r + 1\} \\ &= \#\{\eta \in \mathbb{Z}^2 : 4r \leq |2\eta - 2\delta|^2 < 4(r + 1)\} \\ &= \sum_{l=4r}^{4r+3} \#\{\eta \in \mathbb{Z}^2 : |2\eta - 2\delta|^2 = l\}. \end{aligned}$$

But $|2\eta - 2\delta|^2 = 4|\eta|^2 - 4\langle\eta, 2\delta\rangle + 2 \equiv 2 \pmod{4}$, so the only contribution to the above sum comes from $l = 4r + 2$. Thus, by (10), for any $\varepsilon' > 0$ there exists $c_{\varepsilon'}$ such that

$$(12) \quad \#\{\eta \in \mathbb{Z}^2 : r \leq |\eta - \delta|^2 < r + 1\} \leq c_{\varepsilon'}(4r + 2)^{\varepsilon'} \leq c_{\varepsilon'}(6r)^{\varepsilon'}.$$

Next we observe that for $\delta \in \{(0, \frac{1}{2}), (\frac{1}{2}, 0), (\frac{1}{2}, 1), (1, \frac{1}{2})\}$ we have $|2\eta - 2\delta|^2 \equiv 1 \pmod{4}$, so that the estimate (12) is valid in these cases, too. Iterating the argument, we obtain for $\delta = (\frac{m_1}{2^m}, \frac{m_2}{2^m})$ with $m \in \mathbb{N}$ and $0 \leq m_{1,2} \leq 2^m$ the estimate

$$(13) \quad \#\{\eta \in \mathbb{Z}^2 : r \leq |\eta - \delta|^2 < r + 1\} \leq c_{\varepsilon'}(6^m r)^{\varepsilon'}.$$

Now for an arbitrary $\delta \in [0, 1]^2$ we choose $\delta' = (\frac{m_1}{2^{m'}}, \frac{m_2}{2^{m'}})$ with $|\delta - \delta'| \sim r^{-\frac{1}{2}}$, so that

$$\{\eta \in \mathbb{Z}^2 : r \leq |\eta - \delta|^2 < r + 1\} \subset \{\eta \in \mathbb{Z}^2 : r - 1 \leq |\eta - \delta|^2 < r + 2\}$$

and hence, by (13),

$$\#\{\eta \in \mathbb{Z}^2 : r \leq |\eta - \delta|^2 < r + 1\} \leq 3c_{\varepsilon'}(6^{m'} r)^{\varepsilon'}.$$

Such a δ exists for $2^{m'} \sim r^{\frac{1}{2}}$, estimating roughly, for $6^{m'} \leq r^{\frac{3}{2}}$. So we have the bound

$$\#\{\eta \in \mathbb{Z}^2 : r \leq |\eta - \delta|^2 < r + 1\} \leq 3c_{\varepsilon'} r^{\frac{5\varepsilon'}{2}}.$$

Choosing $\varepsilon' = \frac{2\varepsilon}{5}$, $c_{\varepsilon} = 3c_{\varepsilon'}$, we obtain (11). $\square$

Corollary 1. *If B is a disc (or square) of arbitrary position and of radius (side-length) R , then for any $\varepsilon > 0$ there exists c_{ε} such that*

$$(14) \quad \sum_{\substack{\eta_1 \in \mathbb{Z}^2 \\ r \leq |\eta_1 - \delta|^2 < r+1}} \chi_B(\eta_1) \leq c_{\varepsilon} R^{\varepsilon}.$$

Proof. If $R \gtrsim r^{\frac{1}{6}}$, the estimate (14) follows from Lemma 1. If $R \ll r^{\frac{1}{6}}$, there are at most two lattice points on the intersection of B with the circle of radius $\simeq r^{\frac{1}{2}}$ around δ , by Lemma 4.4 of [4]. $\square$

In the sequel we will use the following projections: For a subset $M \subset \mathbb{Z}^2$ we define P_M by $\mathcal{F}P_M u(k, \eta, \tau) = \chi_M(\eta) \mathcal{F}u(k, \eta, \tau)$. Especially, if M is a ball of radius 2^l centered at the origin, we will write P_l instead of P_M . Furthermore we have $P_{\Delta l} = P_l - P_{l-1}$, and the P -notation will also be used in connection with a sequence $\{Q_{\alpha}^l\}_{\alpha \in \mathbb{Z}^2}$ of squares of sidelength 2^l , centered at $2^l \alpha$. Double sized squares with the same centers will be denoted by $\tilde{Q}_{\alpha}^l$.

Theorem 5. *Let $s > 1$, $b > \frac{1}{2}$ and $\varepsilon > 0$. Then for a disc (or square) B of arbitrary position with radius (sidelength) R we have*

$$(15) \quad \|(P_B u)v\|_{L^2_{xyt}} \lesssim R^{\varepsilon} \|u\|_{X_{0,b}} \|v\|_{X_{s,b}}.$$

Proof. Choose f, g with $\|f\|_{L^2_{\xi\tau}} = \|u\|_{X_{0,b}}$ and $\|g\|_{L^2_{\xi\tau}} = \|v\|_{X_{s,b}}$. Then the left hand side of (15) becomes

$$(16) \quad \left\| \int d\tau_1 \sum_{k_1 \in \mathbb{Z}} \sum_{\eta_1 \in \mathbb{Z}^2} \chi_B(\eta_1) f(\xi_1, \tau_1) \langle \sigma_1 \rangle^{-b} g(\xi_2, \tau_2) |k_2|^{-s} \langle \sigma_2 \rangle^{-b} \right\|_{L^2_{\xi\tau}}.$$

Since $\|uv\|_{L^2_{xyt}} = \|u\bar{v}\|_{L^2_{xyt}}$, which corresponds to $\|\widehat{u}*\widehat{v}\|_{L^2_{\xi\tau}} = \|\widehat{u}*\widehat{v}\|_{L^2_{\xi\tau}}$ on Fourier side, where $\widehat{v}(\xi, \tau) = \widehat{\bar{v}}(-\xi, -\tau)$, and since the phase function ϕ is assumed to be odd, so that $\|u\|_{X_{s,b}} = \|\bar{u}\|_{X_{s,b}}$, we may assume in the estimation on (16), that k_1 and k_2 have the same sign, cf. Remark 4.7 in [2]. So it's sufficient to consider $0 < |k_2| \leq |k_1| < |k|$. Now, using Minkowski's inequality we estimate (16) by

$$\begin{aligned} & \left\| \sum_{k_1 \in \mathbb{Z}} |k_2|^{-s} \int d\tau_1 \sum_{\eta_1 \in \mathbb{Z}^2} \chi_B(\eta_1) f(\xi_1, \tau_1) \langle \sigma_1 \rangle^{-b} g(\xi_2, \tau_2) \langle \sigma_2 \rangle^{-b} \right\|_{L^2_{\eta\tau}} \|L^2_k\| \\ & \leq \| |k_2|^{-\frac{1}{2}} \| \int d\tau_1 \sum_{\eta_1 \in \mathbb{Z}^2} \chi_B(\eta_1) f(\xi_1, \tau_1) \langle \sigma_1 \rangle^{-b} g(\xi_2, \tau_2) \langle \sigma_2 \rangle^{-b} \|_{L^2_{\eta\tau}} \|L^2_{kk_1}\|, \end{aligned}$$

where Cauchy-Schwarz was applied to $\sum_{k_1 \in \mathbb{Z}}$. Thus it is sufficient to show that

$$\begin{aligned} (17) \quad & \left\| \int d\tau_1 \sum_{\eta_1 \in \mathbb{Z}^2} \chi_B(\eta_1) f(\xi_1, \tau_1) \langle \sigma_1 \rangle^{-b} g(\xi_2, \tau_2) \langle \sigma_2 \rangle^{-b} \right\|_{L^2_{\eta\tau}} \\ & \lesssim R^\varepsilon |k_2|^{\frac{1}{2}} \|f(k_1, \cdot, \cdot)\|_{L^2_{\eta_1\tau_1}} \|g(k_2, \cdot, \cdot)\|_{L^2_{\eta\tau}}. \end{aligned}$$

By the "Schwarz-method" developed in [14], [15] and by [5, Lemma 4.2], (17) follows from

$$(18) \quad \sum_{\eta_1 \in \mathbb{Z}^2} \chi_B(\eta_1) \langle \tau - \phi_0(k_1) - \phi_0(k_2) + \frac{|\eta_1|^2}{k_1} + \frac{|\eta_2|^2}{k_2} \rangle^{-2b} \lesssim R^{2\varepsilon} |k_2|.$$

For $\omega := \eta_1 - \frac{k_1}{k}\eta$ we have $\frac{|\eta_1|^2}{k_1} + \frac{|\eta_2|^2}{k_2} = \frac{|\eta|^2}{k} + \frac{k}{k_1 k_2} |\omega|^2$, so that with $a := \tau - \phi_0(k_1) - \phi_0(k_2) + \frac{|\eta|^2}{k}$ the left hand side of (18) becomes

$$\sum_{\eta_1 \in \mathbb{Z}^2} \chi_B(\eta_1) \langle a + \frac{k}{k_1 k_2} |\omega|^2 \rangle^{-2b} = \sum_{r \geq 0} \langle a + \frac{k}{k_1 k_2} r \rangle^{-2b} \sum_{r \leq |\eta_1 - \frac{k_1}{k}\eta|^2 < r+1} \chi_B(\eta_1).$$

By Corollary 1 the inner sum is controlled by $c_\varepsilon R^\varepsilon$, while

$$\sum_{r \geq 0} \langle a + \frac{k}{k_1 k_2} r \rangle^{-2b} \lesssim \frac{|k_1 k_2|}{|k|} \lesssim |k_2|,$$

which proves (18). $\square$

Remark:

The quantity, which we precisely loose in the application of Lemma 1, is

$$r^\varepsilon \simeq |\eta_1 - \frac{k_1}{k}\eta|^{2\varepsilon} \leq \langle k\eta_1 - k_1\eta \rangle^{2\varepsilon},$$

which is the symbol of the Fourier multiplier $M^{2\varepsilon}$. Rereading carefully the calculation in the previous proof, we see that - instead of (17) - the following estimate holds true as well.

$$\begin{aligned} (19) \quad & \left\| \int d\tau_1 \sum_{\eta_1 \in \mathbb{Z}^2} \langle k\eta_1 - k_1\eta \rangle^{-\varepsilon} f(\xi_1, \tau_1) \langle \sigma_1 \rangle^{-b} g(\xi_2, \tau_2) \langle \sigma_2 \rangle^{-b} \right\|_{L^2_{\eta\tau}} \\ & \lesssim \frac{|k_1 k_2|^{\frac{1}{2}}}{|k|^{\frac{1}{2}}} \|f(k_1, \cdot, \cdot)\|_{L^2_{\eta_1\tau_1}} \|g(k_2, \cdot, \cdot)\|_{L^2_{\eta\tau}}. \end{aligned}$$

(Introducing the $M^{-\varepsilon}$ we cannot justify the sign assumption on $k, k_{1,2}$ any more.) Multiplying by $|k|^{\frac{1}{2}}$ and summing up over k_1 using Cauchy-Schwarz we obtain

$$(20) \quad \|\mathcal{F}_x D_x^{\frac{1}{2}} M^{-\varepsilon}(u, v)\|_{L_k^\infty L_{yt}^2} \lesssim \|u\|_{X_{\frac{1}{2},b}} \|v\|_{X_{\frac{1}{2},b}},$$

from which (9) follows by a further application of the Cauchy-Schwarz inequality. So Theorem 2 is proved.

Proof of Theorem 1. Since in Corollary 1 the position of the disc is arbitrary, we may replace $\chi_B(\eta_1)$ by $\chi_B(\eta_2)$ in the proof of Theorem 5, which gives

$$(21) \quad \|u(P_B v)\|_{L_{xyt}^2} \lesssim R^\varepsilon \|u\|_{X_{0,b}} \|v\|_{X_{s,b}}.$$

Now we have symmetry between u and v , so that we may interpolate bilinearly to obtain

$$(22) \quad \|(P_B u)v\|_{L_{xyt}^2} \lesssim R^\varepsilon \|u\|_{X_{s_1,b}} \|v\|_{X_{s_2,b}}$$

for $s_{1,2} \geq 0$ with $s_1 + s_2 > 1$. Decomposing dyadically we obtain with $0 < \varepsilon' < \varepsilon$

$$\begin{aligned} \|uv\|_{L_{xyt}^2} &\leq \sum_{l \geq 0} \|(P_{\Delta l} u)v\|_{L_{xyt}^2} \lesssim \sum_{l \geq 0} 2^{l\varepsilon'} \|P_{\Delta l} u\|_{X_{s_1,b}} \|v\|_{X_{s_2,b}} \\ &\lesssim \sum_{l \geq 0} 2^{l(\varepsilon' - \varepsilon)} \|u\|_{X_{s_1,\varepsilon,b}} \|v\|_{X_{s_2,b}} \lesssim \|u\|_{X_{s_1,\varepsilon,b}} \|v\|_{X_{s_2,b}}. \end{aligned}$$

Exchanging u and v again we have shown for $s_{1,2} \geq 0$ with $s_1 + s_2 > 1$ and $\varepsilon_{1,2} \geq 0$ with $\varepsilon_1 + \varepsilon_2 > 0$ that

$$(23) \quad \|uv\|_{L_{xyt}^2} \lesssim \|u\|_{X_{s_1,\varepsilon_1,b}} \|v\|_{X_{s_2,\varepsilon_2,b}},$$

which is the $\varepsilon_0 = 0$ part of (4) in Theorem 1. To see the $\varepsilon_0 > 0$ part, we decompose

$$\|D_y^{-\varepsilon_0}(uv)\|_{L_{xyt}^2} \leq \sum_{l \geq 0} 2^{-l\varepsilon_0} \|P_{\Delta l}(uv)\|_{L_{xyt}^2},$$

where for fixed l

$$\|P_{\Delta l}(uv)\|_{L_{xyt}^2}^2 = \sum_{\alpha, \beta \in \mathbb{Z}^2} \langle P_{\Delta l}((P_{Q_\alpha^l} u)v), P_{\Delta l}((P_{Q_\beta^l} u)v) \rangle_{L_{xyt}^2}.$$

Now for $\eta_1 \in Q_\alpha^l$, $|\eta| \leq 2^l$ we have $\eta_2 = \eta - \eta_1 \in \tilde{Q}_{-\alpha}^l$, so that the latter can be estimated by

$$\begin{aligned} &\sum_{\alpha, \beta \in \mathbb{Z}^2} \langle (P_{Q_\alpha^l} u)(P_{\tilde{Q}_{-\alpha}^l} v), (P_{Q_\beta^l} u)(P_{\tilde{Q}_{-\beta}^l} v) \rangle_{L_{xyt}^2} \\ &\leq \sum_{\alpha, \beta \in \mathbb{Z}^2} \langle (P_{\tilde{Q}_\alpha^l} u)(P_{\tilde{Q}_\beta^l} \bar{v}), (P_{\tilde{Q}_\beta^l} u)(P_{\tilde{Q}_\alpha^l} \bar{v}) \rangle_{L_{xyt}^2} \\ &\leq \sum_{\alpha, \beta \in \mathbb{Z}^2} \| (P_{\tilde{Q}_\alpha^l} u)(P_{\tilde{Q}_{-\beta}^l} v) \|_{L_{xyt}^2} \| (P_{\tilde{Q}_\beta^l} u)(P_{\tilde{Q}_{-\alpha}^l} v) \|_{L_{xyt}^2} \\ &\leq \sum_{\alpha, \beta \in \mathbb{Z}^2} \| (P_{\tilde{Q}_\alpha^l} u)(P_{\tilde{Q}_{-\beta}^l} v) \|_{L_{xyt}^2}^2. \end{aligned}$$

Using (22) and the almost orthogonality of the sequence $\{P_{\tilde{Q}_\alpha^l} v\}_{\alpha \in \mathbb{Z}^2}$ we estimate the latter by

$$2^{2l\varepsilon} \sum_{\alpha, \beta \in \mathbb{Z}^2} \|P_{\tilde{Q}_\alpha^l} u\|_{X_{s_1,b}}^2 \|P_{\tilde{Q}_{-\beta}^l} v\|_{X_{s_2,b}}^2 \lesssim 2^{2l\varepsilon} \|u\|_{X_{s_1,b}}^2 \|v\|_{X_{s_2,b}}^2.$$

Choosing $\varepsilon < \varepsilon_0$ the sum over l remains finite and we arrive at

$$\|D_y^{-\varepsilon_0}(uv)\|_{L_{xyt}^2} \lesssim \|u\|_{X_{s_1,b}} \|v\|_{X_{s_2,b}}.$$

Finally we remark that (4) and (5) are equivalent by duality. $\square$

3. APPLICATIONS TO KP-II TYPE EQUATIONS

Here the phase function is specified as $\phi_0(k) = |k|^\alpha k$, $\alpha \geq 2$, so that the mixed weight becomes $\sigma = \tau - |k|^\alpha k + \frac{|\eta|^2}{k}$. To prove the well-posedness results in Theorem 3 and 4, we need some more norms and function spaces, respectively. In both cases we use the spaces $X_{s,\varepsilon,b;\beta}$ with additional weights, introduced in [2] and defined by

$$(24) \quad \|f\|_{X_{s,\varepsilon,b;\beta}} := \left\| \langle k \rangle^s \langle \eta \rangle^\varepsilon \langle \sigma \rangle^b \left(1 + \frac{\langle \sigma \rangle}{\langle k \rangle^{\alpha+1}} \right)^\beta \hat{f} \right\|_{L_{\tau\xi}^2},$$

We will always have $\beta \geq 0$, so that

$$(25) \quad \|f\|_{X_{s,b}} \leq \|f\|_{X_{s,b;\beta}}.$$

Observe that

$$(26) \quad \|f\|_{X_{s,b}} \sim \|f\|_{X_{s,b;\beta}},$$

if $\langle \sigma \rangle \leq \langle k \rangle^{\alpha+1}$.

The case $\alpha = 2$ corresponding to the original KP-II equation becomes a limiting case in our considerations, where we have to choose the parameter $b = \frac{1}{2}$. Thus we also need the auxiliary norms

$$(27) \quad \|f\|_{Y_{s,\varepsilon;\beta}} := \left\| \langle k \rangle^s \langle \eta \rangle^\varepsilon \langle \sigma \rangle^{-1} \left(1 + \frac{\langle \sigma \rangle}{\langle k \rangle^{\alpha+1}} \right)^\beta \hat{f} \right\|_{L_\xi^2(L_\tau^1)},$$

cf. [5], and

$$(28) \quad \|f\|_{Z_{s,\varepsilon;\beta}} := \|f\|_{Y_{s,\varepsilon;\beta}} + \|f\|_{X_{s,\varepsilon,-\frac{1}{2};\beta}}.$$

As before, for $\varepsilon = 0$ we will write $X_{s,b;\beta}$ instead of $X_{s,\varepsilon,b;\beta}$, and if the exponent β of the additional weight is zero, we use $X_{s,\varepsilon,b}$ as abbreviation for $X_{s,\varepsilon,b;\beta}$. Similar for the Y - and Z -norms. In these terms the crucial bilinear estimate leading to Theorem 3 is the following.

Lemma 2. *Let $\alpha = 2$, $s \geq \frac{1}{2}$ and $\varepsilon > 0$. Then there exists $\gamma > 0$, such that for all u, v supported in $[-T, T] \times \mathbb{T}^3$ the estimate*

$$(29) \quad \|\partial_x(uv)\|_{Z_{s,\varepsilon;\frac{1}{2}}} \lesssim T^\gamma \|u\|_{X_{s,\varepsilon,\frac{1}{2};\frac{1}{2}}} \|v\|_{X_{s,\varepsilon,\frac{1}{2};\frac{1}{2}}}$$

holds true.

Correspondingly for Theorem 4 we have

Lemma 3. *Let $3 < \alpha \leq 4$. Then, for $s > \frac{3-\alpha}{2}$ there exist $b' > -\frac{1}{2}$ and $\beta \in [0, -b']$, such that for all $b > \frac{1}{2}$*

$$(30) \quad \|D_x^{s+1+\varepsilon} M^{-\varepsilon}(u, v)\|_{X_{0,b';\beta}} \lesssim \|u\|_{X_{s,b;\beta}} \|v\|_{X_{s,b;\beta}},$$

whenever $\varepsilon > 0$ is sufficiently small, and

$$(31) \quad \|\partial_x(uv)\|_{X_{s,b';\beta}} \lesssim \|u\|_{X_{s,b;\beta}} \|v\|_{X_{s,b;\beta}}.$$

In the proof of both Lemmas above the resonance relation for the KP-II-type equation with quadratic nonlinearity plays an important role. We have

$$(32) \quad \sigma_1 + \sigma_2 - \sigma = r(k, k_1) + \frac{|k\eta_1 - k_1\eta|^2}{kk_1k_2},$$

where

$$|r(k, k_1)| = ||k|^\alpha k - |k_1|^\alpha k_1 - |k_2|^\alpha k_2| \sim |k_{max}|^\alpha |k_{min}|,$$

see [9]. Both terms on the right of (32) have the same sign, so that

$$(33) \quad \max\{|\sigma|, |\sigma_1|, |\sigma_2|\} \gtrsim |k_{min}| |k_{max}|^\alpha + \frac{|k\eta_1 - k_1\eta|^2}{|kk_1k_2|}.$$

The proof of Lemma 2 is almost the same as that of Lemma 4 in [7], it is repeated here - with minor modifications - for the sake of completeness. We need a variant of Theorem 1 with $b < \frac{1}{2}$. To obtain this, we first observe that, if $s_{1,2} \geq 0$ with $s_1 + s_2 > \frac{1}{2}$, $\varepsilon_{0,1,2} \geq 0$ with $\varepsilon_0 + \varepsilon_1 + \varepsilon_2 > 1$, $1 \leq p \leq 2$, and $b > \frac{1}{2p}$, then

$$(34) \quad \|\mathcal{F}D_y^{-\varepsilon_0}(uv)\|_{L_\xi^2 L_\tau^p} \lesssim \|u\|_{X_{s_1, \varepsilon_1, b}} \|v\|_{X_{s_2, \varepsilon_2, b}}.$$

This follows from Sobolev type embeddings and applications of Young's inequality. Dualizing the $p = 2$ part of (34) we obtain

$$(35) \quad \|uv\|_{X_{-s_1, -\varepsilon_1, -b}} \lesssim \|D_y^{\varepsilon_0} u\|_{L_{xyt}^2} \|v\|_{X_{s_2, \varepsilon_2, b}}.$$

Now bilinear interpolation with Theorem 1 gives the following.

Corollary 2. *Let $s_{1,2} \geq 0$ with $s_1 + s_2 = 1$ and $\varepsilon_{1,2} \geq 0$ with $\varepsilon_1 + \varepsilon_2 > 0$, then there exist $b < \frac{1}{2}$ and $p < 2$ such that*

$$(36) \quad \|\mathcal{F}(uv)\|_{L_\xi^2 L_\tau^p} + \|uv\|_{L_{xyt}^2} \lesssim \|u\|_{X_{s_1, \varepsilon_1, b}} \|v\|_{X_{s_2, \varepsilon_2, b}}$$

and

$$(37) \quad \|uv\|_{X_{-s_1, -b}} \lesssim \|D_y^{\varepsilon_1} u\|_{L_{xyt}^2} \|v\|_{X_{s_2, \varepsilon_2, b}}$$

hold true.

The purpose of the $p < 2$ part in the above Corollary is to deal with the Y -contribution to the Z -norm in Lemma 2. Its application will usually follow on an embedding

$$\|\langle \sigma \rangle^{-\frac{1}{2}} \widehat{f}\|_{L_\xi^2 L_\tau^1} \lesssim \|\widehat{f}\|_{L_\xi^2 L_\tau^p},$$

where $p < 2$ but arbitrarily closed to 2. Now we're prepared to establish Lemma 2.

Proof of Lemma 2. Without loss of generality we may assume that $s = \frac{1}{2}$. The proof consists of the following case by case discussion.

Case a: $\langle k \rangle^3 \leq \langle \sigma \rangle$. First we observe that

$$(38) \quad \|\partial_x(uv)\|_{Z_{s, \varepsilon; \frac{1}{2}}} \lesssim \|D_x^{s+1}(D_y^\varepsilon u \cdot v)\|_{Z_{0,0; \frac{1}{2}}} + \|D_x^{s+1}(u \cdot D_y^\varepsilon v)\|_{Z_{0,0; \frac{1}{2}}}.$$

The first contribution to (38) equals

$$\begin{aligned} & \|\mathcal{F}(D_y^\varepsilon u \cdot v)\|_{L_{\xi, \tau}^2} + \|\langle \sigma \rangle^{-\frac{1}{2}} \mathcal{F}(D_y^\varepsilon u \cdot v)\|_{L_\xi^2 L_\tau^1} \\ & \lesssim \|\mathcal{F}(D_y^\varepsilon u \cdot v)\|_{L_{\xi, \tau}^2 \cap L_\xi^2 L_\tau^p} \lesssim \|u\|_{X_{s, \varepsilon, b}} \|v\|_{X_{s, \varepsilon, b}} \end{aligned}$$

by (36), for some $b < \frac{1}{2}$. Using the fact¹ that under the support assumption on u the inequality

$$(39) \quad \|u\|_{X_{s, \varepsilon, b}} \lesssim T^{\tilde{b}-b} \|u\|_{X_{s, \varepsilon, \tilde{b}}}$$

¹for a proof see e. g. Lemma 1.10 in [6]

holds, whenever $-\frac{1}{2} < b < \tilde{b} < \frac{1}{2}$, this can, for some $\gamma > 0$, be further estimated by $T^\gamma \|u\|_{X_{s,\varepsilon,\frac{1}{2},\frac{1}{2}}} \|v\|_{X_{s,\varepsilon,\frac{1}{2},\frac{1}{2}}}$ as desired. The second contribution to (38) can be treated in precisely the same manner.

Case b: $\langle k \rangle^3 \geq \langle \sigma \rangle$. Here the additional weight on the left is of size one, so that we have to show

$$\|\partial_x(uv)\|_{Z_{s,\varepsilon}} \lesssim T^\gamma \|u\|_{X_{s,\varepsilon,\frac{1}{2},\frac{1}{2}}} \|v\|_{X_{s,\varepsilon,\frac{1}{2},\frac{1}{2}}}.$$

Subcase b.a: σ maximal. Exploiting the resonance relation (33), we see that the contribution from this subcase is bounded by

$$\|\mathcal{F} D_x D_y^\varepsilon (D_x^{-\frac{1}{2}} u \cdot D_x^{-\frac{1}{2}} v)\|_{L_{\xi,\tau}^2 \cap L_{\xi}^2 L_{\tau}^p} \lesssim \|\mathcal{F} (D_x^{\frac{1}{2}} D_y^\varepsilon u \cdot D_x^{-\frac{1}{2}} v)\|_{L_{\xi,\tau}^2 \cap L_{\xi}^2 L_{\tau}^p} + \dots,$$

where $p < 2$. The dots stand for the other possible distributions of derivatives on the two factors, in the same norms, which - by (36) of Corollary 2 - can all be estimated by $c \|u\|_{X_{s,\varepsilon,b}} \|v\|_{X_{s,\varepsilon,b}}$ for some $b < \frac{1}{2}$. The latter is then further treated as in case a.

Subcase b.b: σ_1 maximal. Here we start with the observation that by Cauchy-Schwarz and (39), for every $b' > -\frac{1}{2}$ there is a $\gamma > 0$ such that

$$\|\partial_x(uv)\|_{Z_{s,\varepsilon}} \lesssim T^\gamma \|D_x^{s+1}(uv)\|_{X_{0,\varepsilon,b'}}.$$

With the notation $\Lambda^b = \mathcal{F}^{-1} \langle \sigma \rangle^b \mathcal{F}$ we obtain from the resonance relation that

$$\begin{aligned} \|D_x^{s+1}(uv)\|_{X_{0,\varepsilon,b'}} &\lesssim \|D_x(D_x^{-\frac{1}{2}} \Lambda^{\frac{1}{2}} u \cdot D_x^{-\frac{1}{2}} v)\|_{X_{0,\varepsilon,b'}} \\ &\lesssim \|(D_x^{\frac{1}{2}} D_y^\varepsilon \Lambda^{\frac{1}{2}} u)(D_x^{-\frac{1}{2}} v)\|_{X_{0,b'}} + \|(D_x^{\frac{1}{2}} \Lambda^{\frac{1}{2}} u)(D_x^{-\frac{1}{2}} D_y^\varepsilon v)\|_{X_{0,b'}} \\ &\quad + \|(D_x^{-\frac{1}{2}} D_y^\varepsilon \Lambda^{\frac{1}{2}} u)(D_x^{\frac{1}{2}} v)\|_{X_{0,b'}} + \|(D_x^{-\frac{1}{2}} \Lambda^{\frac{1}{2}} u)(D_x^{\frac{1}{2}} D_y^\varepsilon v)\|_{X_{0,b'}} \end{aligned}$$

Using (37) the first two contributions can be estimated by $c \|u\|_{X_{s,\varepsilon,\frac{1}{2},\frac{1}{2}}} \|v\|_{X_{s,\varepsilon,b}}$ as desired. The third and fourth term only appear in the frequency range $|k| \ll |k_1| \sim |k_2|$, where the additional weight in the $\|u\|_{X_{s,\varepsilon,\frac{1}{2},\frac{1}{2}}}$ -norm on the right becomes $\frac{|k_2|}{|k_1|}$, thus shifting a whole derivative from the high frequency factor v to the low frequency factor u . So, using (37) again, these contributions can be estimated by

$$c \|u\|_{X_{s,\varepsilon,\frac{1}{2},\frac{1}{2}}} \|v\|_{X_{s,\varepsilon,b}} \lesssim \|u\|_{X_{s,\varepsilon,\frac{1}{2},\frac{1}{2}}} \|v\|_{X_{s,\varepsilon,b;\frac{1}{2}}}.$$

□

Now we turn to the proof of Lemma 3, where the restrictions to the b -parameters can be relaxed slightly, so that the auxiliary Y - and Z -norms are not needed. We use again the Λ -notation, i. e. $\Lambda^b = \mathcal{F}^{-1} \langle \sigma \rangle^b \mathcal{F}$.

Proof of Lemma 3. First we show how (30) implies (31). By the resonance relation (33) we have

$$|k_1 \eta - k \eta_1|^2 \lesssim |k k_1 k_2| (\langle \sigma \rangle + \langle \sigma_1 \rangle + \langle \sigma_2 \rangle) \leq |k k_1 k_2| \langle \sigma \rangle \langle \sigma_1 \rangle \langle \sigma_2 \rangle,$$

so that (31) is reduced to

$$\|D_x^{s+1+\frac{\varepsilon}{2}} M^{-\varepsilon}(u, v)\|_{X_{0,b'+\frac{\varepsilon}{2};\beta}} \lesssim \|u\|_{X_{s-\frac{\varepsilon}{2},b-\frac{\varepsilon}{2};\beta}} \|v\|_{X_{s-\frac{\varepsilon}{2},b-\frac{\varepsilon}{2};\beta}},$$

Relabelling appropriately and choosing ε sufficiently small, we see that (31) follows from (30). To prove the latter, we may assume $s \leq 0$. Next we choose ε small and b' close to $-\frac{1}{2}$ so that

$$(40) \quad s > 2 + (\alpha + 1)b' + 3\varepsilon$$

and $\beta := \frac{s-b'}{\alpha} \in [0, -b']$. Now the proof consists again of a case by case discussion.

Case a: $\langle k \rangle^{\alpha+1} \leq \langle \sigma \rangle$. Here it is sufficient to show

$$(41) \quad \|D_x^{s+1+\varepsilon-\alpha\beta-\beta} M^{-\varepsilon}(u, v)\|_{X_{0,b'+\beta}} \lesssim \|u\|_{X_{s,b}} \|v\|_{X_{s,b}}$$

Subcase a.a: $|k| \ll |k_1| \sim |k_2|$.

Subsubcase triple a: $\langle \sigma \rangle \geq \langle \sigma_{1,2} \rangle$. Here we use the resonance relation (33) to see that the left hand side of (41) is bounded by

$$\begin{aligned} & \|D_x^{s+1+\varepsilon-\alpha\beta+b'} M^{-\varepsilon}(D_x^{\frac{\alpha(b'+\beta)}{2}} u, D_x^{\frac{\alpha(b'+\beta)}{2}} v)\|_{L_{xyt}^2} \\ & \lesssim \|M^{-\varepsilon}(D_x^{\frac{s+1+\varepsilon+(\alpha+1)b'}{2}} u, D_x^{\frac{s+1+\varepsilon+(\alpha+1)b'}{2}} v)\|_{L_{xyt}^2}, \end{aligned}$$

where we have used the assumption on the frequency sizes in this subcase. Observe that our choice of β implies $s+1+\varepsilon-\alpha\beta+b' = 1+\varepsilon+2b' \geq 0$. Now the bilinear estimate (9) is applied to obtain the upper bound

$$\|D_x^{\frac{s+2+3\varepsilon+(\alpha+1)b'}{2}} u\|_{X_{0,b}} \|D_x^{\frac{s+2+3\varepsilon+(\alpha+1)b'}{2}} v\|_{X_{0,b}} \lesssim \|u\|_{X_{s,b}} \|v\|_{X_{s,b}},$$

where in the last step we have used (40).

Subsubcase a.a.b: $\langle \sigma_1 \rangle \geq \langle \sigma \rangle, \langle \sigma_2 \rangle$. Here the resonance relation (33) gives that the left hand side of (41) is bounded by

$$\begin{aligned} & \|D_x^{s+1+\varepsilon-\alpha\beta+b'} M^{-\varepsilon}(D_x^s \Lambda^b u, D_x^{\alpha(b'+\beta)-s} v)\|_{X_{0,-b}} \\ & \lesssim \|D_x^{-\frac{1}{2}-\varepsilon} M^{-\varepsilon}(D_x^s \Lambda^b u, D_x^{\frac{3}{2}+2\varepsilon+(\alpha+1)b'} v)\|_{X_{0,-b}}. \end{aligned}$$

Now the dual version of estimate (9), that is

$$(42) \quad \|M^{-\varepsilon}(u, v)\|_{X_{-\frac{1}{2}-, -\frac{1}{2}-}} \lesssim \|u\|_{L_{xyt}^2} \|v\|_{X_{\frac{1}{2}+, \frac{1}{2}+}}$$

is applied, which gives, together with the assumption (40), that the latter is bounded by $c\|u\|_{X_{s,b}} \|v\|_{X_{s,b}}$. This completes the discussion of subcase a.a. Concerning subcase a.b, where $|k| \gtrsim |k_{1,2}|$, we solely remark that it can be reduced to the estimation in subsubcase triple a.

Case b: $\langle k \rangle^{\alpha+1} \geq \langle \sigma \rangle$. Here the additional weight in the norm on the left of (30) is of size one, so our task is to show

$$(43) \quad \|D_x^{s+1+\varepsilon} M^{-\varepsilon}(u, v)\|_{X_{0,b'}} \lesssim \|u\|_{X_{s,b;\beta}} \|v\|_{X_{s,b;\beta}}.$$

Subcase b.a: $\langle \sigma \rangle \geq \langle \sigma_{1,2} \rangle$. Here we may assume by symmetry that $|k_1| \geq |k_2|$. We apply (33) and (9) to see that for $\delta \geq 0$ the left hand side of (43) is controlled by

$$\begin{aligned} & \|M^{-\varepsilon}(D_x^{s+1+\varepsilon+\alpha b'+\delta} u, D_x^{b'-\delta} v)\|_{L_{xyt}^2} \\ & \lesssim \|D_x^{\frac{3}{2}+2\varepsilon+\alpha b'+\delta} u\|_{X_{s,b}} \|D_x^{b'+\frac{1}{2}+\varepsilon-\delta} v\|_{X_{0,b}}. \end{aligned}$$

The latter is bounded by $c\|u\|_{X_{s,b}} \|v\|_{X_{s,b}}$, provided $\frac{3}{2} + 2\varepsilon + \alpha b' + \delta \leq 0$ and $b' + \frac{1}{2} + \varepsilon - \delta \leq s$, which can be fulfilled by a proper choice of $\delta \geq 0$, since (40) holds.

Subcase b.b: $\langle \sigma_1 \rangle \geq \langle \sigma \rangle, \langle \sigma_2 \rangle$.

Subsubcase b.b.a: $|k_1| \gtrsim |k_2|$. With $\delta \geq 0$ as in subcase b.a the contribution here is bounded by

$$\begin{aligned} & \|D_x^{-\frac{1}{2}-\varepsilon} M^{-\varepsilon} (D_x^{s+\frac{3}{2}+2\varepsilon+\alpha b'+\delta} \Lambda^b u, D_x^{b'-\delta} v)\|_{X_{0,-b}} \\ & \lesssim \|D_x^{\frac{3}{2}+2\varepsilon+\alpha b'+\delta} u\|_{X_{s,b}} \|D_x^{b'+\frac{1}{2}+\varepsilon-\delta} v\|_{X_{0,b}} \lesssim \|u\|_{X_{s,b}} \|v\|_{X_{s,b}}, \end{aligned}$$

where (33) and the dual version (42) of Theorem 2 were used again. Finally we turn to the

Subsubcase triple b, where $|k_1| \ll |k| \sim |k_2|$. Here the additional weight in $\|u\|_{X_{s,b;\beta}}$ on the right of (43) behaves like

$$\left(\frac{|k|}{|k_1|}\right)^{\alpha\beta} \sim \left(\frac{|k_2|}{|k_1|}\right)^{\alpha\beta},$$

so that it is sufficient to show

$$(44) \quad \|D_x^{s+1+\varepsilon} M^{-\varepsilon} (u, D_x^{-\alpha\beta} v)\|_{X_{0,b'}} \lesssim \|u\|_{X_{s-\alpha\beta,b}} \|v\|_{X_{s,b}}.$$

Now by (33) the left hand side of (44) can be controlled by

$$\begin{aligned} & \|D_x^{-\frac{1}{2}-\varepsilon} M^{-\varepsilon} (D_x^{b'} \Lambda^b u, D_x^{s+\frac{3}{2}+2\varepsilon+\alpha(b'-\beta)} v)\|_{X_{0,-b}} \\ & \lesssim \|D_x^{b'} u\|_{X_{0,b}} \|D_x^{2+3\varepsilon+(\alpha+1)b'} v\|_{X_{0,b}} \end{aligned}$$

by (42). Since $b' = s - \alpha\beta$ the first factor equals $\|u\|_{X_{s-\alpha\beta,b}}$, while by (40) the second is dominated by $\|v\|_{X_{s,b}}$. This proves (44). $\square$

Finally we recall the definition of the Fourier restriction norm spaces from [2]. For a time slab $I = (-\delta, \delta) \times \mathbb{T}^3$ they are given by

$$X_{s,\varepsilon,b;\beta}^\delta := \{u|_I : u \in X_{s,\varepsilon,b;\beta}\}$$

with norm

$$\|u\|_{X_{s,\varepsilon,b;\beta}^\delta} := \inf\{\|\tilde{u}\|_{X_{s,\varepsilon,b;\beta}} : \tilde{u} \in X_{s,\varepsilon,b;\beta}, \tilde{u}|_I = u\}.$$

Now our well-posedness results read as follows.

Theorem 6 (precise version of Theorem 3). *Let $s \geq \frac{1}{2}$ and $\varepsilon > 0$. Then for $u_0 \in H_x^s H_y^\varepsilon(\mathbb{T}^3)$ satisfying (2) there exist $\delta = \delta(\|u_0\|_{H_x^s H_y^\varepsilon}) > 0$ and a unique solution $u \in X_{s,\varepsilon,\frac{1}{2};\frac{1}{2}}^\delta$ of the Cauchy problem (1) with $\alpha = 2$. This solution is persistent and the mapping $u_0 \mapsto u$, $H_x^s H_y^\varepsilon(\mathbb{T}^3) \rightarrow X_{s,\varepsilon,\frac{1}{2};\frac{1}{2}}^{\delta_0}$ is locally Lipschitz for any $\delta_0 \in (0, \delta)$.*

Theorem 7 (precise version of Theorem 4). *Let $3 < \alpha \leq 4$, $s \geq s' > \frac{3-\alpha}{2}$ and $\varepsilon \geq 0$. Then for $u_0 \in H_x^s H_y^\varepsilon(\mathbb{T}^3)$ satisfying (2) there exist $b > \frac{1}{2}$, $\beta > 0$, $\delta = \delta(\|u_0\|_{H_x^{s'} H_y^\varepsilon}) > 0$ and a unique solution $u \in X_{s,\varepsilon,b;\beta}^\delta \subset C^0((-\delta, \delta), H_x^s H_y^\varepsilon(\mathbb{T}^3))$. This solution depends continuously on the data, and extends globally in time, if $s \geq 0$ and $\varepsilon = 0$.*

With the estimates from Lemma 2 and 3 at our disposal the proof of these theorems is done by the contraction mapping principle, cf. [2], [5], [14], [15]. The reader is also referred to section 1.3 of [6], where the related arguments are gathered in a general local well-posedness theorem.

REFERENCES

- [1] Ben-Artzi, M., Saut, J.-C.: Uniform decay estimates for a class of oscillatory integrals and applications, *Differential and Integral Equations* 12 (1999), no. 2, 137–145.
- [2] Bourgain, J.: On the Cauchy problem for the Kadomtsev-Petviashvili equation, *Geom. Funct. Anal.* 3 (1993), no. 4, 315–341.
- [3] Bourgain, J.: Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. I. Schrödinger equations, *GAFA* 3 (1993), 107 - 156.
- [4] De Silva, D., Pavlovic, N., Staffilani, G., Tzirakis, N.: Global Well-Posedness for a periodic nonlinear Schrödinger equation in 1D and 2D. *Discrete Contin. Dyn. Syst.* 19 (2007), 37 - 65
- [5] Ginibre J., Tsutsumi, Y., Velo G.: On the Cauchy problem for the Zakharov system. *J. Funct. Anal.* 151 (1997), 384–436.
- [6] Grünrock, A.: New applications of the Fourier restriction norm method to wellposedness problems for nonlinear evolution equations. Thesis, Universität Wuppertal, 2002
- [7] Grünrock, A., Panthee, M., Silva, J.: On KP-II type equations on cylinders. Preprint, 2008.
- [8] Hadac, M.: On the local well-posedness of the Kadomtsev-Petviashvili II equation. Thesis, Universität Dortmund, 2007
- [9] Hadac, M.: Well-posedness for the Kadomtsev-Petviashvili equation (KP-II) and generalisations, *Trans. Amer. Math. Soc.*, S 0002-9947(08)04515-7, 2008.
- [10] Hadac, M., Herr, S., Koch, H.: Well-posedness and scattering for the KP-II equation in a critical space. *Annales Henri Poincaré* (to appear).
- [11] Hardy, G.H., Wright, E.M.: An introduction to the theory of numbers. 5th ed. Oxford etc.: Oxford at the Clarendon Press. XVI, 1979
- [12] Isaza, P., Lopez, J., Mejia, J.: The Cauchy problem for the Kadomtsev-Petviashvili (KP-II) equation in three space dimensions, *Comm. Partial Differential Equations*, 32 (2007), 611–641.
- [13] Isaza, P., Mejia, J.: Local and global Cauchy problems for the Kadomtsev-Petviashvili (KP-II) equation in Sobolev spaces of negative indices, *Comm. Partial Differential Equations*, 26 (2001), 1027–1054.
- [14] Kenig, C., Ponce, G., Vega, L.: A bilinear estimate with applications to the KdV equation, *J. of the AMS* 9 (1996), 573 - 603
- [15] Kenig, C., Ponce, G., Vega, L.: Quadratic forms for the 1 - D semilinear Schrödinger equation, *Transactions of the AMS* 348 (1996), 3323 - 3353
- [16] Saut, J.-C.: Remarks on the generalized Kadomtsev-Petviashvili equations, *Indiana Univ. Math. J.* 42 (1993), no. 3, 1011–1026.
- [17] Saut, J.-C., Tzvetkov, N.: The Cauchy problem for higher-order KP equations, *J. Differential Equations* 153 (1999), 196–222.
- [18] Saut, J.-C., Tzvetkov, N.: The Cauchy problem for the fifth order KP equations, *J. Math. Pures Appl.* 79, 4 (2000), 307–338.
- [19] Saut, J.-C., Tzvetkov, N.: On Periodic KP-I Type Equations, *Commun. Math. Phys.* 221 (2001), 451–476.
- [20] Takaoka, H., Tzvetkov, N.: On the local regularity of the Kadomtsev-Petviashvili-II equation, *IMRN* (2001), no. 2, 77–114.
- [21] Takaoka, H., Tzvetkov, N.: On 2D nonlinear Schrödinger equations with data on $\mathbb{R} \times \mathbb{T}$. *J. Funct. Anal.* 182 (2001), 427 -442
- [22] Tzvetkov, N.: On the Cauchy problem for Kadomtsev Petviashvili equation. *Comm. PDE*, 24 (1999), 1367-1397

AXEL GRÜNROCK: RHEINISCHE FRIEDRICH-WILHELMS-UNIVERSITÄT BONN, MATHEMATISCHES INSTITUT, BERLINGSTRASSE 1, 53115 BONN, GERMANY.

E-mail address: gruenroc@math.uni-bonn.de